

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/320696573>

Taxonomy and morphological study on the vertebrate remains of Shark and rays fauna from the Middle and Late Eocene succession, Fayoum Depression, Egypt

Article · January 2017

CITATIONS
0

READS
479

4 authors, including:



Abdelfattah Zalata
Tanta University

70 PUBLICATIONS 291 CITATIONS

[SEE PROFILE](#)



Hamza Moustafa Khalil
Tanta University

30 PUBLICATIONS 197 CITATIONS

[SEE PROFILE](#)



Mohamed Sobhy FATHY
Tanta University

13 PUBLICATIONS 80 CITATIONS

[SEE PROFILE](#)


Some of the authors of this publication are also working on these related projects:



Diatom nanotechnology [View project](#)



Implementation of the Geosciences to Construct the New Desert Urban, Site Management, and Distribute Resources; Pilot area: Moghra Oasis, Qattara Depression. [View project](#)



Taxonomy and morphological study on the vertebrate remains of Shark and rays fauna from the Middle and Late Eocene succession, Fayoum Depression, Egypt

Abdelfattah A. Zalata, Hamza M. Khalil, Mohamed S. Fathy and Rana M. Tarek

Department of Geology, Fac. of Science, Tanta University, Tanta, Egypt
(E-mail: abzalata@science.tanta.edu.eg)

Abstract: Well preserved vertebrate remains of Shark and ray fauna from the Middle and Late Eocene succession of Fayoum depression, at Gebal Qasr El-Sagha area have been documented and studied in details. Four stratigraphic sections are measured, described and sampled in the field at Hussein Wally Village, Birket Qarun, Qasr el-Sagha and Wadi el-Afreet. Lithostratigraphically, the studied succession is divided into three formations arranged from base to top into Gehannam, Birket Qarun and Qasr el-Sagha formations. This Eocene sequence provides by far the most complete view of the endemic African vertebrate fauna. Identification of the basic pattern of fish remains and taxonomic evaluation revealed that the recorded shark and rays taxa belong to 3 classes, 7 orders, 12 families, 18 genera, and 21 species. The identified taxa are macro-scale, collected on the surface, and known either from teeth or rostral remains. A taxonomic account and detailed morphologic description of the fossils shark, rays and bony fish teeth have been achieved. The depositional environments in the studied Middle-Late Eocene age sequence are interpreted. The abundance of recognized vertebrate fauna indicates environments varying from open marine shelf with low energy conditions to restricted marine shallow water conditions. However, the frequent distribution of macrofauna with intense bioturbation in sandstones of Birket Qarun Formation is a good indicator of restricted shallow water conditions.

Key words: Taxonomy, Vertebrates remains, Eocene, Stratigraphy, Fayoum depression

Introduction:

The Fayoum depression was subjected to many geological and archaeological studies over 200 years, as it holds a rich heritage of best paleontological and archeological sites in the world. Among these studies (Said et al. 1972; Butzer 1976; Wendorf and Schild 1976; Kozłowski 1983; Hassan 1985, 1986, 1997; Said 1990; Hendrickx and Vermeersch 2000). The depression hosts a unique heritage in terms of vertebrates and mammals fossils. It has been known as a repository of Paleogene vertebrates and has attracted the attention of scientists worldwide (e.g., Kirk and Simons 2001; Simons 2005; Lewis and Simons 2007; Simons et al. 2007; Antar 2011; Adnet et al. 2011; Underwood et al. 2011). The Middle and Late Eocene represented an important period in the evolution of sharks and rays and saw the establishment of 'modern-type' trophic systems (Underwood et al. 2011).

The Eocene rocks around and to the west of the Fayoum depression comprise a thick succession of shelf marine rocks representing environments from open shelf to restricted lagoon. The stratigraphy of the area has been documented by several authors (e.g. Beadnell 1905; Said 1962; Iskander 1943; Allam et al. 1991; Gingerich 1992; Shama and Shided 1994; Abdullah et al. 1997; Dolson et

al. 2002; Ismail and Abd El-Azeam 2008; Abu El Ghar 2012). However, a considerable lateral variation within parts of the succession has caused problems with applying a lithostratigraphical scheme (Strougo 2008).

The Fayoum depression has a fundamentally important role to play in developing an understanding of Egyptian prehistory, in particular during the Epipalaeolithic – Neolithic time. This depression has been inhabited by humans as early as 8500 BP. The first significant occupation by humans, which is documented archaeologically, occurred during the epipalaeolithic and Neolithic Age, around 8200 BP. (Hassan 1986). During this time clear traces of human impact on the Fayoum depression could be observed in the old Fayoum lake sediments (Zalata 2015).

Description of study area

The Fayoum depression occupies nearly triangular depression, located between latitudes 29° 00' – 29° 45' N and longitudes 30° 00' – 31° 10' E, immediately to the west of the Nile Valley and about 90 km to the south west of Cairo (Fig.1). The depression covers an area of approximately 1700 km² and it lies below sea level. The Fayoum is separated from the Nile Valley in the east by a

ridge running south from the Giza plateau. The ridge is approximately 8-10 km wide at its northern boundary, but narrows to 2.5km in the south. Its surface is flat and slopes downward northwesterly from 32 m above mean sea level to 45 m below mean sea level at Qarun Lake, which is the lowest part of the depression. The Fayoum depression is carved out of the Middle- Upper Eocene bedrock of the Western Desert of Egypt, and is surrounded by escarpments from all sides. The Eocene sequence is differentiated into four rock units including Wadi El-Rayan, Gehannam, Birket Qarun and Qasr El-Sagha Formations.

Material and Methods

Four stratigraphic sections are measured, described and sampled in the studied area. The first section (I) is located in south west Qarun Lake at Hussein Wally Village, between Longitude 29°27' 56"N and Latitude 30°23'36.8" E. The second section (II) is obtained from the cliffs bordering the south western shore of lake Qarun and far about 4Km east from the first section, between longitude 29°27'47.1" N and Latitude 30°26'1.0" E. The third section (III) is measured at Qasr El-Sagha Temple, between Longitude 29°36'3.2" N and Latitude 30°40'13.9" E. The last section (IV) is located far about 3.5 Km N-E Qasr El-Sagha Temple (Wadi El-Afreet Section), between Longitude 29°37'10.3" N and Latitude 30°41'54.8" E. A total of 130 samples are collected from the studied sections and investigated in details for paleontological and microfacies analyses.

Lithostratigraphy

The general lithostratigraphic units for Middle-Upper Eocene sequences proposed by Beadnell (1905) and modified later by Said (1962) are the most suitable and will be followed herein. The Umm Rigl Member of Gingerich (1992) is used herein to define the lowermost strata of the Qasr El-Sagha Formation in the Hussein Walley Village and Birket Qarun sections. Moreover, the Temple and the Dir Abu Lifa members of Qasr El-Sagha Formation suggested by Bown and Kraus (1988) in the vicinity of Qasr El-Sagha Temple are adopted. Three rock units are recognized as follows from the base to the top; Gehannam, Birket Qarun and Qasr El-Sagha Formations. The last formation is included Umm Rigl, Temple and Dir Abu Lifa members.

Gehannam Formation

Author: Said (1962)

Type locality and Type section: The Garret Gehannam, south west Fayoum Province. It is contained about 35m of gypsiferous claystone, marly limestone, marly sandstone, glauconitic sandstone and marls (Said 1962).

Measured sections: The base of the formation is not exposed, but it is likely that the lowest sample was from a level quite high in the formation. About 15 to 20m of this formation are exposed in the base of Hussein Walley Village and Birket Qarun sections respectively.

Boundaries: In the west Fayoum area, the Gehannam Formation conformably overlies El Gharag Formation and underlies the Birket Qarun Formation.

Description: The exposed section of upper part Gehannam Formation in the west south Qarun Lake is distinguished mainly by gypsiferous calcareous claystones and shale with marl intercalations (Fig. 3). The beds are bioturbated, sometimes glauconitic and contain a diverse, open marine fauna. The upper part of the Gehannam Formation passes upwardly into the mudstones and sandstones of the Birket Qarun Formation.

Birket Qarun Formation

Author: Beadnell (1905)

Type locality and type section: It is represented in cliffs bordering the northern shore of lake Qarun (Said 1962), and the type-section is thus the steep-faced precipitous escarpment described by Beadnell (1905).

Measured sections: This formation is well developed in the two studied sections at south-western end of Lake Qarun. Its thickness of about 38m at Hussein Wally Village section and increase to about 52.5m at Birket Qarun section (Fig. 3).

Boundaries: The Birket Qarun Formation conformably overlies the Gehannam Formation by stratigraphic gradual contact and underlies the Qasr El Sagha Formation.

Description: In the studied sections, the Birket Qarun Formation consists mainly of mudstone, siltstones and, heavily bioturbated calcareous sandstones, and trace fossil banks at its top part. The lower part of this formation is arranged in coarsening-upward sequences starting with claystones, siltstones and ending with fine to medium sized sandstone. This sandstone characterizes by golden yellow color, carbonate concretions, calcareous, fossiliferous thin beds with trace fossils in parts. The size and numbers of carbonate concretions increase upwardly and to north direction in the studied area. The upper part of the Formation consists mainly of brownish yellow sandstone with some shale layer intercalations. The shale change in thickness laterally and characterizes by *Thalassinoides* trace fossils. On other hand the top most sandstone bed at the contact with superimposed Qasr El-Sagha Formation is highly bioturbated by *Rhizoliths* trace fossils.

Qasr El Sagha Formation

Author: Beadnell (1905)

Type locality and Type section: It is located at Qasr El Sagha Temple with a thickness 175 m of alternating series of clays and limestone with sands and sandstone in the upper beds.

Measured sections: The Qasr El Sagha Formation is recorded in all studied sections and distinguished by three members. Um Rigl Member, the lower part of the Qasr El Sagha Formation attains 51m at Birket Qarun section and 31m at Hussein Walley Village section. In addition to the two upper members (Temple and Dir Abu Lifa) are measured at Qasr El Sagha and Wadi El Afreet sections by 88m and 95 respectively (Fig. 3).

Boundaries: In the studied area, the Qasr El Sagha Formation conformably overlies the Birket Qarun Formation and conformably and/or unconformably underlies the Gabel Qatrani Formation.

Description and subdivisions: The Qasr El Sagha Formation consists mainly of a thick succession of dark gypsiferous mudstones with thin hash fossiliferous reddish bed intercalations, and overlain by friable, cleaned washed, cross bedded sandstones and ends with an extensively hard yellow carbonate layer known as the Bare Limestone (very hard dolomitic limestone).

Results and Systematic description

The recorded vertebrate remains from the studied Eocene successions are investigated in detail. The results of vertebrate identification show 21 species belong to 18 genera, 12 families, 7 orders and 3 classes. The recognized fish assemblage has clear ecological affinities with Eocene Tethyan fauna and also has common elements with Eocene myliobatids, Carcharhinidae, Triakidae and Hemigaleidae fishes. The most common taxa are included *Hexanchus agassizi* Cappetta, *Pristis lathami* Galeotti, *Otodussokolovi* (Jaekel), *Macrorhizodus praecursor* (Leriche), *Cretolamna twiggsensis* (Case), *Galeocерdo latidens* (Agassiz), *Negaprion frequens* (Dames) *Rhinoptera sherburni* Arambourg, *Carcharodon carcharias* Linnaeus, *Galeocерdo cuvieri* Péron & Lesueur 1822, *Xiphiorhynchus aegyptiacus* Weiler, *Rhynchobatus* sp., *Carcharias* sp., *Negaprion* sp., *Aetobatus* sp., *Galeocерdo* sp., *Rhizoprionodon* sp., *Leidybatis* sp., and *Eutrachiurides* sp.

The classification followed here after Compagno et al. 2005

- Class Chondrichthyes Huxley 1880
- Subclass Elasmobranchii Bonaparte 1836
- Cohort Euselachii Hay 1902
- Subcohort Neoselachii Compagno 1977
- Super order Squalomorphii Compagno 1973
- Order Hexanchiformes Buen 1926

Suborder Hexanchoidei Garman 1913

Family Hexanchidae Gray 1851

Genus *Hexanchus* Rafinesque 1810

Hexanchus agassizi Cappetta 1976

Pl. 1, fig. 15.

2013 *Hexanchus agassizi*, Otero et al., Fig. 5: 1-6

2014 *Hexanchus* cf. *agassizi*, Carlsen & Cuny, Fig. 16 E-F

Description: The best preserved tooth measures 11 mm mesio-distally, 6 mm apico-basally and 1.7 mm labio-lingually. The tooth is worn. It is comb-shaped with a main cusp and seven accessory cusps regularly decreasing in size distally. All cusps are leaning apicodistally. The mesial cutting edge of the main cusp has no serration and no heel. It is convex in the basal two thirds and straight near the apex. The distal cutting edge is concave in the basal part and convex in the apical part. The lingual face is convex, the labial face almost flat.

Geological distribution: The genus is known since the Early Jurassic to Recent (Cappetta, 1987), with a cosmopolitan distribution. Ward (1979) describes the teeth of *Hexanchus agassizi*, from the Lower Eocene of England. *Hexanchus agassizi* is known from southwestern France and from the London Clay of England (Adnet 2006). Carlsen & Cuny Fig (2014) 16E-F (Early-Middle Eocene Denmark). Ypresian of England, Eocene of New Jersey, and the Oligocene of Australia and the the Lower Oligocene of ex-U.S.S.R. (Cappetta, 1987). Eocene *Hexanchus* has been described from Seymour Island, Antarctica (Cione and Reguero, 1994).

Known geologic range: Cretaceous – Oligocene (Otero et al. 2012)

Hexanchus sp.

2006 *Hexanchus* sp. Adnet, Fig. 3

2006 *Hexanchus griseus* Adnet, Fig. 2 (A-L),

Remarks: Small teeth, with very distinct cutting edges and wide crown. The recorded taxon is similar to *Hexanchus* sp. and *Hexanchus griseus* (Bonnaterre, 1788) that reported by Adnet (2006) from late Ypresian/early Lutetian, south-western France.

Order: Rajiformes Berg, 1940

Suborder Pristioidei CAPPETTA 1980

Family: Pristidae BONAPARTE 1838

Genus *Pristis* Latham, 1794

Pristis lathami Galeotti, 1837

Pl. 1, fig. 18

2012 *Pristis lathami* Galeotti, Diedrich, Fig. 14 (18).

2012 *Pristis lathami* Galeotti, Zalmout et al., Fig. 50-P

Description: The tooth is 45 mm long, 15 mm wide and 7 mm thick at the base of the crown. The proximal half of the tooth is thickest, and the tooth tapers distally to a much thinner 3 mm. The anterior edge of the tooth is thick and blunt, while the posterior edge is concave, a shallow gutter that runs from the base of the tooth to the

distal end. The vertebrae is rounded and has concentric lines in internal view and longitudinal lines in side view.

Geographic distribution: *Pristis lathami* is known from lower and upper Eocene deposits in Africa and North America (Cappetta, 1987). Eocene of the Paris Basin (Casier, 1949). In Egypt the species is known from the Eocene of the Fayoum and Bahariya regions (Stromer, 1905B; Case and Cappetta, 1990; Underwood *et al.*, 2011; Murray *et al.*, 2011; Adnet *et al.*, 2011). Late Eocene Qattara depression, Egypt (Zalmout *et al.* 2012), late Eocene of South Carolina and North America (Cicimurri 2007), Southern North Sea (Diedrich 2012). Farther east, *Pristis lathami* is known from the Eocene of Qatar (Casier, 1971). Middle–Late Eocene, southwestern Morocco (Adnet *et al.* 2010)

Known Geologic range: Early to Late Eocene.

***Pristis aquitanicus* Delfortrie 1872**

1959 *Pristis aquitanicus*, Delfortrie, Ghosh, p. 67, pl. 88, Figs. 7, 8.

2013 *Pristis aquitanicus* Delfortrie, Sharma and Patnaik, Fig. 5: J, K.

Description: The anterior surface is tapering gently towards the tip thereby increasing the curvature while the posterior surface is straight and has a broad sulcus. Length of about 22 mm and width 21 mm.

Geographic distribution: The fossil remains of the genus *Pristis* ranging in age from Lower Eocene to Recent (Sharma and Patnaik 2013). It is reported from Miocene of India (Sharma and Patnaik 2013). Cappetta (1970) also reported *P. aquitanicus* from the Middle Miocene of Southern France

Known Geologic range: Early to Late Eocene.

Suborder Rhynchobatoidei Flower 1941

Family: Rhynchobatidae Garman, 1913

Genus: *Rhynchobatus* Müller and Henle, 1837

***Rhynchobatus* sp.**

2010 *Rhynchobatus* sp. Adnet *et al.*, Fig. 4 E.

2011 *Rhynchobatus* sp. Underwood *et al.*, Fig. 7 (X).

2011 *Rhynchobatus* sp. Antar Fig. 29 (O)

2013 *Rhynchobatus* sp. Sharma and Patnaik, Fig. 5: F.

Description: A small tooth; crown enamel is having a granular texture, the lingual face is flat, concave on each side separated by a moderate medial uvula, the terminal end of the median uvula is pointed, lateral uvulae absent; root massive and extends beyond the lingual face of the crown; root lobes divided on both the sides by a deep nutritive groove and a small foramina.

Geographic distribution: It is recorded from Middle to Late Eocene of the Fayoum area (Underwood *et al.*, 2011), Late Eocene of the western Desert, Egypt (Adnet *et al.*, 2010), Langhian of Loupian, Herault, Southern

France (Cappetta, 1987) and from Oligocene Chandler Bridge Formation of South Carolina, U.S. (Cicimurri and Knight, 2009). *Rhynchobatus* teeth have also been reported from the Maastrichtian of Morocco (Arambourg, 1952), the Miocene of Japan (Itoigawa *et al.*, 1985) and Miocene of India (Sharma and Patnaik, 2013).

Known geologic range: Maastrichtian to Miocene.

Suborder Myliobatoidei Stingrays

Super family Myliobatoidea Compango 1973

Family: Myliobatoidea Bonaparte 1838

Genus: *Myliobatis* Cuvier 1817

***Myliobatis* sp.**

1972 *Myliobatis* sp., Welton, Plate 1, p. 167, Fig. 8.

2011 *Myliobatis* sp., Antar, Fig 29 J-J1

2011 *Myliobatis* sp., Underwood *et al.*, Fig. 4 (G).

2012 *Myliobatis* sp. Otero *et al.*, Fig. 3: T

2013 *Myliobatis* sp., Sharma and Patnaik, Fig. 3(A).

Description: Teeth broader than length, with hexagonal contour in shape and rectilinear outline; the length is about 28 mm and the width of about 7-9 mm, the crown is as thick as the root, flat and quite convex, being displaced anteriorly with respect to the root. The crown and the root are well separated by a minor groove. The basal surface of the root is flat with 23 root lobes separated by grooves. The tooth is much broader than long.

Geographic distribution: The genus *Myliobatis* is known since the Early Paleocene to Recent, with a cosmopolitan distribution (Cappetta, 1987). *Myliobatis* sp. has been described from the Paleocene to Eocene of western Africa including Angola, Egypt, Morocco, Togo and Nigeria (Cappetta 1987; Cook *et al.* 2010). Middle Eocene of Kutch (Mishra, 1980), Miocene Coaledo Formation along the Oregon coast Welton (1972), Miocene of Baripada Beds, Orissa (Sahni and Mehrotra, 1981), Eocene of Subathu Formation, Bilaspur area and Himachal Pradesh (Singh, 1985, Kumar and Loyal, 1987), Eocene Vastan lignite Mine of Cambay Shale, Gujarat (Rana *et al.*, 2004), Lower Eocene Panandro lignite field, Gujarat (Bajpai *et al.*, 2002), Early Eocene Kapurdi Formation of Rajasthan (Rana *et al.*, 2006), Eocene deposits of Fayoum and western Desert, Egypt (Case and Cappetta, 1990; Antar, 2011; Underwood *et al.*, 2011), Miocene of India (Sharma and Patnaik, 2013). The genus also has been recognized on Seymour Island, Antarctica (Kriwet, 2005).

Known geologic range: Campanian to Recent.

Genus: *Aetobatus* Blainville 1816

***Aetobatus* sp.**

2011 *Aetobatus* sp. Antar, Fig. 29 (M).as

2011 *Aetobatus* sp. Underwood Fig. 4 (F)

2013 *Aetobatus* sp. Sharma and Patnaik Fig. 4: E, F, G

Description: One sample is collected and well preserved. Root is quite distinctive with small vertical lines along it and has smooth edges. The crown lacks the hexagonal shape which is usually associated with myliobatids. The contact between root and crown is straight. Crown is slightly serrate and less thickness than root. The occlusal surface is smooth. The root is nearly as high as the crown and is divided into longitudinal ridges and grooves. The length is about 6mm and width of about 6mm.

Geographic distribution: *Aetobatus* teeth have been described from the Lower Miocene of the western coast of India (Sahni and Mishra, 1975).

Known geologic range: Middle Eocene to Lower Miocene

Genus: *Leidybatis* Cappetta, 1986

Leidybatis sp.

Pl.1, fig.17

Description: Teeth broad with rectangular shape. The length is about 28-32 mm and the width of about 7-9 mm, Root is thick and distinctive with small serration along it and has smooth edges. The contact between root and crown is straight. Crown is slightly serrate and less thickness than root. The root is nearly as high as the crown and is divided into longitudinal ridges and grooves.

Known geologic range: Middle to Late Eocene

Order: Myliobatiformes Compagno, 1973

Family: Rhinopteridae Jordan and Evermann, 1896

Genus: *Rhinoptera* Cuvier, 1829

Type species: *Myliobatis marginata* Saint-Hillaire, 1817, Recent, Mediterranean Sea.

Remarks: The Rhinopteridae are comprised of a single genus, *Rhinoptera* (Cappetta 2004)..

Geographic distribution: The genus is known from the Paleocene in North and West Africa, and Europe (Cappetta 2004), and Lower Miocene of Switzerland and France and most other marine Neogene deposits (Leriche 1927; Cappetta 1970).

***Rhinoptera sherburni* Arambourg 1952**

Pl.1, fig.16

1952 *Rhinoptera* aff. *sherburni* Arambourg, Plate XXXII, Figs. 15-24.

2013 *Rhinoptera sherburni* Arambourg, Sharma and Patnaik (Fig. 4: H, I)

Description: The teeth are medium size, with 8-10mm length, and 4-6 mm width, hexagonal in shape; teeth possess a prominent but thin lingual shelf; labial face is more upright, the lingual root overhangs the crown by a distinct margin, the crown is thick. Upper surface of root is slightly curved and smooth. The contact between root and crown is very slightly curved. The crown is serrate and curved in middle area and become straight towards edges.

Geographic distribution: *Rhinoptera sherborni* has been reported from the early Eocene of Virginia (Kent, 1999); the Middle Eocene of England (Kemp et al., 1990), Eocene Nigeria (White, 1926), Morocco (Arambourg, 1952) and Uzbekistan (as *Rhinoptera* cf. *sherborni*, Case et al., 1996) and the late Eocene of Egypt (Murray et al., 2011). Fossil record of *Rhinoptera* extends back upto the Late Palaeocene (Cappetta, 1987, 2006), Middle-Late Eocene, southwestern Morocco (Adnet et al. 2010), Miocene of India (Sharma and Patnaik 2013), Madagascar (Wallett, 2006).

Known geologic range: Paleocene to Miocene

Order Lamniformes BERG 1958

Family: Odontaspidae Müller and Henle, 1839

Genus: *Carcharias* Rafinesque, 1810

***Carcharias* sp.**

2012 *Carcharias* sp. Zalmout et al. Fig. 3B-E

2012 *Carcharias* sp. Diedrich Fig. 14 (7-8)

2013 *Carcharias* sp. Otero et al., Fig. 3: 1-4

Description: The recorded taxon is small-sized not larger than 2 cm, with length 15-18 mm, and width 5-8mm. The crown sharp having smooth enamel in both labial and lingual surfaces; the cutting edge is complete and reaches the base of the crown; slender root branches with medial groove and deep depression between them; In some specimens the root of tooth is partially broken and arched. The contact between root and crown is arched. Crown is thin and long and has smooth edges and surface in lingual and labial view are ended with very acute end.

Geographic distribution: Fossils of *Carcharias* have been found all over the world, especially in the Cretaceous, Eocene, Miocene and Oligocene sediments of Europe, United States, Australia and Africa (Cappetta and Case, 1975a,b, Cappetta, 1987, Otero et al., 2013, Zalmout et al., 2012, Diedrich 2012). Southern North Sea (Diedrich 2012), Late Eocene Qasr El Sgha, Fayoum (Underwood et al. 2011), Late Eocene of Qattara depression, Egypt (Zalmout et al. 2012), Pliocene Farol das Lagostas locality, Angola (Antunes 1978).

Known geologic range: Maastrichtian to Pliocene.

Family Otodontidae Gückman 1964

Genus *Otodus* Jordan and Hannibal, 1923

***Otodus sokolovi* (Jaekel, 1895)**

Pl.1, figs.1-4

2010 *Carcharocles sokolovi* Jaekel, Adnet et al. Fig. 3 a1,2

2011 *Carcharocles sokolovi* Jaekel, Antar, Fig. 24

2011 *Otodus* (*Carcharocles*) *sokolovi* Jaekel, Underwood et al. Fig. 4 A

2012 *Otodus* cf. *sokolovi* Jaekel, Zalmout et al. 2012, Fig. 3F-BB

2012 *Carcharocles sokolovi* Jaekel, Diedrich, Fig. 10 (9-10)

Description: Large teeth that may reach 60 to 80 mm in height and 45-65 mm in width. Central blades are large

and bear fine and regular strong serrations along both cutting edges. The teeth have two divergent lateral cusps which are triangular in shape, serrated and wide.

Geographic Distribution: This species is recorded from middle Eocene of Syria, Jordan (Cappetta et al., 2000, Mustafa and Zalmout, 2002; Smadi et al., 2003; Mustafa et al., 2005), middle Eocene of Fayoum, Egypt (Stromer, 1905A; Case and Cappetta, 1990; Vliet and Abu el Khair, 2010; Underwood et al., 2011), Middle–Late Eocene, southwestern Morocco (Adnet et al., 2010), and Nigeria, Congo, and Angola (White, 1926; Dartevelle and Casier, 1942, 1943, 1949, 1959), Late Eocene Qattara depression (Zalmout et al. 2012), In North America *Otodus* (*Carcharocles*) is known from North Carolina (Bourdon and Chandler, 2007), the Barnwell Formation in Georgia (Case and Borodin, 2000), and middle Eocene of Alabama (White, 1956), middle and late Eocene Tethyan deposits in North Africa, the Middle East, and North America (Zalmout et al. 2012).

Known geologic range: Middle Eocene - Lower Oligocene.

Family: Lamnidae Müller & Henle 1838

Genus *Carcharodon* Müller & Henle, 1838

***Carcharodon carcharias* Linnaeus, 1758**

Pl.1, fig. 9

1973 *Carcharodon carcharias* Linnaeus, Mehrotra et al., p. 191-192, pl. 2, figs. 2, 6 a-b.

1981 *Carcharodon carcharias* Linnaeus, Sahni and Mehrotra, p. 112-113.

1998 *Carcharodon carcharias* Linnaeus, Tiwari et al., p. 12, pl. 1, fig. 5, 6.

2009 *Carcharodon carcharias* Linnaeus, Mondal et al., pl. II, figs. 1-4.

2010 *Carcharodon carcharias* Linnaeus, Andreev and Motchurova, Fig. 2 a, b

Description: The tooth is having height equal or greater than the width. Length 22 mm and width 16 mm. Tooth composed of root and crown. Root is broken except small part and appeared slightly curved and crown is broad in upper part and become narrow towards lower part and has sharp end and serrate edges and has longitudinal lines in lingual and labial view. Serrations of the cutting edges are uniform, and relatively coarser towards the base in larger specimens. Mesial edge is more concave than distal edge, mesial concavity shows wide variations, some are feebly concave or straight. Labial surface is flat, however at the tip it may be slightly inwardly curved. The base of the root may be feebly bifurcated or flat.

Geographic distribution: This species is recorded from Farol das Lagostas, Angola. (Andreev and Motchurova 2010), Miocene Chesapeake Bay at Calvert Cliffs in Maryland, USA, Lowery et al. (2011), Early Pliocene of Spain, Early – Middle Miocene of Greens mill run, north

carollina, western and Eastern USA, South Africa, Australia and Late Eocene Qasr El-Sagha Formation Egypt (Zalmout et al. 2012).

Known geologic Range: Eocene to Early Pliocene.

Genus *Macrorhizodus* Glikman, 1964

Type species: *Isurus praecursor* (Leriche, 1905).

***Macrorhizodus praecursor* (Leriche, 1905)**

Pl.1, fig.5

2011 *Cosmopolitodus praecursor* (*Macrorhizodus praecursor*), Leriche, Antar, Fig.25: A-B7.

2012 *Macrorhizodus praecursor* Leriche, Zalmout et al., Fig. 4A-V

2012 *Isurus praecursor* Leriche, Diedrich, Fig.11 (1-10).

2012 *Macrorhizodus (Isurus) praecursor* Leriche, Otero et al., Fig. 3: L, M

2013 *Macrorhizodus praecursor* Leriche, Otero et al., Fig. 3: 28-34

Description: Medium to relatively large sized lamniform teeth with high, slender and triangular crown that becomes broader toward the base, being labio-lingually compressed; flat labial face with soft enamel and smooth folds near the root; lingual face is convex with some cracks in the enamel; root is slightly massive, basally flattened. The teeth may reach 40-44 mm in total height anteriorly, and 30-32 mm in total width laterally.

Geographic distribution: *Macrorhizodus praecursor* is widely known from all middle and late Eocene Tethyan deposits of Europe, North America and some late Eocene marine vertebrate localities in Egypt (Adnet et al., 2011), Middle to Late Eocene of the Seymour Island in Antarctica (Cione and Reguero, 1994), Priabonian of the southernmost Chile (Otero et al., 2012b). Eocene deposits in Belgium, Syria, Nigeria, Togo, Guinea Bissau and England (Cappetta, 1987). Middle-Late Eocene of Chesapeake Bay, U.S. (Kent, 1994), southwestern Morocco (Adnet et al. 2010) and Southern North Sea (Diedrich (2012). Late Eocene of Qattara depression, Egypt (Zalmout et al., 2012).

Known Geologic range: Middle - late Eocene

Family: Cretoxyrhinidae Gluckman 1958

Genus: *Cretolamna* GLUCKMAN 1958

***Cretolamna twiggensis* Case, 1981**

Pl.1, figs.11-12

1981 *Lamna twiggensis* Case, pl. 3, Figs. 3–8

1990 *Cretolamna twiggensis* Case, Case & Cappetta, pl. 3, Figs. 40–55

2007 *Cretolamna twiggensis* Case, Adnet et al., Figs. 6.21 and 6.22

2011 *Brachycarcharias twiggensis* Case, Underwood et al., Fig. 4L–M

2011 *Cretolamna twiggensis* Case, Adnet et al, Fig. 3A

2012 *Brachycarcharias twiggensis* Case, Zalmout et al., Fig. 4W–X

Description: Teeth are relatively large; with height up to 20-25mm and 15-18 mm width, the cusp is high, quite triangular in upper teeth and not very thick. The root lobes are elongated and their ends are often rounded. The upper anterior tooth has two divergent lobes with a height less than the crown, the anterior –lateral teeth have vertical central blade, slightly sigmoidal in shape and had a wide base, with pointed edge and having two small diverted lateral cusps.

Geographic distribution: the genus is nearly worldwide, Early Cretaceous-Early Eocene. African distribution and occurrence: Morocco, Angola, Democratic Republic of the Congo, Niger, Nigeria and Egypt; Late Cretaceous-Eocene (Cappetta 1987). The species extends to palaeotropical seas between the Caribbean, western Tethys (Case, 1981; Case & Borodin, 2000) and oriental Tethys (Casier, 1971; Case & Cappetta, 1990; Case & West, 1991; Adnet *et al.* 2007). It is recorded from Egypt (Mokkatom to Wadi Hitan areas), both in Bartonian and Priabonian deposits (Case and Cappetta, 1990 and Underwood *et al.*, 2011), middle and late Eocene of the Fayoum and Qattara depression, Western Desert of Egypt (Case and Cappetta 1990; Strougo *et al.*, 2007; Underwood *et al.*, 2011; Adnet *et al.*, 2011; Zalmout *et al.* 2012), the species is recorded elsewhere from Pakistan (Adnet *et al.*, 2007), Middle–Late Eocene of southwestern Morocco (Adnet *et al.*, 2010) and Georgia, USA (Case, 1981; Case and Borodin, 2000).

Known geologic range: Middle–Late Eocene

Order Carcharhiniformes

Family: Carcharhinidae Jordan & Evermann, 1896

Genus *Galeocerdo* Müller and Henle, 1837

Type species: *Galeocerdo cuvieri* Peron and Le Sueur, 1822

Galeocerdo cuvieri Péron & Lesueur, 1822

Pl.1, fig. 14

1973 *Galeocerdo cuvieri* Péron & Lesueur, Mehrotra *et al.*, p. 184, pl. 1, fig. 3.

1981 *Galeocerdo cuvieri* Péron & Lesueur, Sahni and Mehrotra, p. 109, pl.2, fig. 12.

2009 *Galeocerdo cuvieri* Péron & Lesueur, Mondal *et al.*, Pl.I, figs. 11-12.

Description: Tooth large with 23-30 mm long and 18-21 mm width, sub-triangular, irregular in shape, broader than high and highly oblique. Crown is labio-lingually thickened; the mesial cutting edge has a deep notch, distal margin strongly convex; both cutting margins are serrated; strength of serrations becomes weak to obsolete near the apex; at distal cutting edge strength of serrations are coarser at the middle, decreases towards the base and the apex, whereas mesial edge serrations are finer and uniform. Mesial heel is extended, with numerous denticles, size of the denticles increases proximally. On the labial surface a prominent triangular pit is present at the middle near the root; labial surface bears few longitudinal striations; crown-root boundaries at both

faces are convex, convexity more pronounced in lingual side. Root higher than crown; in cross-section it is labio-lingually arched and thicker at the middle part; in profile the tooth deflects outward.

Geographic distribution: *G. cuvieri* has been reported from the lower Miocene beds of different localities in Gujarat (Mehrotra *et al.* 1973). Miocene from baripada, Orissa (Mondal *et al.* 2009). The type species, *Galeocerdo cuvieri* (Peron and Le Sueur 1822), is extant and can be found in all tropical and temperate seas, including those of Madagascar (Smale 1998; Cappetta 2004). Pliocene of Italy (Lawley 1876), South Africa (Davies 1964), and North Carolina (Cappetta 2004).

Known geologic range: Middle Eocene to Pliocene.

Galeocerdo sp.

1990 *Galeocerdo* sp. Case and Cappetta, 1990, plate 5, figs 92-95.

2010 *Galeocerdo* sp. Andreev and Motchurova, fig. 5 a-c; fig. 6

2002 *Galeocerdo* sp. Bajpai and Thewissen, txt-fig.2f.

Description: Root of the tooth is arched and elongate and has linear groove in mid line in lingual view and smooth in labial view. The contact between root and crown is slightly arched.

Geographic distribution: Late Eocene of El-Sagha Formation (Case and Cappetta, 1990). Teeth of *Galeocerdo* have previously been reported in Egypt by Stromer (1905a) and Priem (1897b). Eocene of Minqar Tabaghbagh, western Desert, Egypt (Vliet and Abu El-Khair 2010). This species is recorded from Upper Miocene of Angola. (Andreev and Motchurova 2010).

Known Geologic range: Eocene to Upper Miocene.

Genus *Carcharhinus* Blainville, 1816

Type Species: *Carcharias melanopterus* Quoy and Gaimard, 1824

Carcharhinus sp.

2010 *Carcharhinus* sp. Andreev P, Motchurova, Fig. 5 c, d

2010 *Carcharhinus* sp. Adnet *et al.* Fig.3 G.

2011 *Carcharhinus* sp. 1, Antar, Fig. 28 C

2012 *Carcharhinus* sp.1. Zalmout *et al.*, Fig. 5E-F.

Description: The teeth are relatively small up to 11 mm height and 11-12mm width. Cusp is flattened labially and triangular in shape, extends mesiodistally into shoulders; the shoulders are sharp and separated from the main blade by slightly developed notches. Lingually, it is convex and the apex curved labially and has no lateral cusplets. The roots have a shallow furrow in the middle of the lingual portion.

Geographic distribution: Middle–Late Eocene, southwestern Morocco (Adnet *et al.* 2010), Middle and

Late Eocene deposits of Fayoum and Qattara depressions, Egypt (Underwood et al. 2011, Zalmout et al., 2012)

Known Geologic range: Middle and Late Eocene.

Genus *Negaprion* Whitley, 1940

***Negaprion frequens* (Dames, 1883)**

Pl.1, fig.10

1908 *Carcharias* (*Aprionodon*) aff. *frequens* Priem, pl.15, Fig. 6–7

1971 *Aprionodon frequens* Casier, pl. 1, Fig. 6

1990 *Carcharhinus frequens* Dames, Case & Cappetta, pl. 5, Figs. 102– 107; pl.7, Figs. 143–148 and 151–159

1990 *Negaprion frequens* Dames, Case and Cappetta, Plate 7, 147–147.

2011 *Carcharhinus* aff. *frequens* Dames, Adnet et al., Fig. 3G–H

2011 *Negaprion frequens* Dames, Underwood et al., Fig. 5V

2012 *Negaprion frequens*, Dames, Zalmout et al., Fig. 5, A–D

Description: Teeth of are relatively small, ranging from 7 to 12 mm in total height, from 5 to 12 mm in total width (mesiodistally), and from 2 to 3 mm in lingolabial thickness in the middle of the root. The upper teeth have a crown of triangular outline with a cusp slightly slanted distally, while the lower teeth generally have a slender cusp with smooth cutting edges than never reach the heels, except on some lateral teeth. The cutting edges are always unserrated and are not disconnected from the cutting edges of lateral heels which are totally smooth. The root lobes have enlarged and rounded ends.

Geographic distribution: It is recorded from the Miocene of Europe (Leriche 1926; Leriche 1957; Antunes and Jonet 1970; Cappetta 1970), North Africa (Arambourg 1952), North and South America (Longbottom 1979), and India (Mehrotra et al 1973; Sahni and Mehrotra 1981), Angola (Antunes 1978), middle Eocene of the Midra and Saila shales of Qatar (Casier, 1971), Late Eocene of Birket Qarun and Qasr el-Sagha formations, Fayoum area (Case and Cappetta, 1990; Underwood et al., 2011), late Eocene of Jordan (Cappetta et al., 2000; Mustafa and Zalmout, 2002), and the late middle Eocene to late Eocene of southwestern Morocco (Adnet et al., 2010), Miocene fishes from baripada beds, Orissa (Patnaik et al., 2014).

Known geologic range: Eocene to Pleistocene.

Genus: *Rhizoprionodon* Whitley 1929

Type species: *Carcharias* (*Scoliodon*) *crenidens* Klunzinger, 1880.

***Rhizoprionodon* sp.**

Pl.1, figs. 7–8

1990 *Rhizoprionodon* sp., Case & Cappetta, pl.7. Figs. 160–163

1991 *Rhizoprionodon* sp., Case & West, pl. 3. Figs. 2–4

2011 *Rhizoprionodon* sp., Adnet et al., Fig. 3P–Q

2012 *Rhizoprionodon* sp., Zalmout et al., Fig. 5K

2013 *Rhizoprionodon* sp., Otero et al., Fig. 6: 3–8

2014 *Rhizoprionodon* sp., Sharma and Patnaik, Pl. 4, figs. 9, 10

Description: Teeth are small, up to 7–8mm height and 8–10mm width, and 2–3mm thick, cusp small and is bent backward with extended crown at the base. Both the cutting edges are sharp, smooth without any serration. The mesial cutting edge is long slightly concave and recurved towards the apex and the distal edge is short; the mesial cutting edge is sigmoidal. The lingual face is convex and the labial face is flat. The distal heels are rounded with cusplet. The crown overhangs the root. The root low and broad, median nutritive groove is present in the lingual face of the root. The basal margin of the root concave to straight.

Geographic distribution: This genus is relatively common in worldwide Eocene marine deposits, Fayoum and Qattara depression, Egypt (Case and Cappetta, 1990; Strougo et al., 2007; Underwood et al., 2011), from Pakistan (Case and West, 1991; Adnet et al., 2007), Middle to Late Eocene. Río Baguales Formation (Otero et al. 2012). *Rhizoprionodon* sp. is recorded from the Eastern Coast of India (Patnaik et al., 2014).

Known geologic range: Middle Eocene to Miocene.

Class: Osteichthyes Huxley, 1880

Order: Perciformes Bleeker, 1859

Family: Trichiuridae Rafinesque, 1810

Genus: *Eutrichiurides* Casier 1944

***Eutrichiurides* sp.**

Pl.1, fig.13

1966 *Eutrichiurides* sp. pl.3, fig. 30

2004 *Eutrichiurides* sp. Rana et al., Fig. 3 (41)

Description: Teeth long and slender with conical shape; height up to 33 mm and width of about 8 mm. Apex small chisel-like, about one-tenth of basal part; slight keel along the edges extend towards base; basal part laterally compressed with vertical ridges and grooves. It has irregular surface and longitudinal lines on lingual and dorsal view.

Geographic distribution: *Eutrichiurides* is an extinct genus of prehistoric bony fish. This genus is recorded from the Cretaceous age at the Khouribga Plateau, Morocco, the Early-Mid Eocene, Monmouth County, the Early Eocene (Ypresian) of the lower part of the Jaisalmer basin (Cappetta 1987).

Known geologic range: Cretaceous -Eocene

Class Actinopterygii (sensu Nelson, 1994)

Division Teleostei Muller, 1844

Order Perciformes Bleeker, 1859

Suborder Xiphioidei Swainson, 1839

Family Xiphiidae Swainson, 1839

Subfamily Xiphiorhynchinae Regan, 1909

Genus *Xiphiorhynchus* van Beneden, 1871

Type Species-*Xiphiorhynchus elegans* van Beneden, 1871

***Xiphiorhynchus aegyptiacus* Weiler, 1929**

1929 *Xiphiorhynchus aegyptiacus*. Weiler, Taf. I, Fig. 4.

Description: Elongate vertebrae, thin thickness in the middle that become broad towards lower and upper edges and has knobs in labial view. Height of about 12-15 mm and width 8-13 mm. In cross section, the rostrum contains two types of longitudinal canals, an unpaired central canal and two pairs of lateral nutrient canals, and both types vary as to how far they extend distally. The dorsal pair of lateral nutrient canals is positioned closer to the mid-line than the ventral pair of lateral nutrient canals.

Known Geologic range: Eocene

Summary and conclusion

The studied Eocene succession exposed at the northern part of Lake Qarun is differentiated into three rock units arranged from base to top: Gehannam, Birket Qarun and Qasr El-Sagha formations. Qasr El-Sagha Formation is distinguished into three members: Um Reigl, Temple and Abu Lifa members. Paleontologically, the vertebrate faunal content of the studied sections is investigated. A total of 21 sharks species are recorded and suggest that they occupied a wide range of ecological niches. Some of the recorded species are limited in their stratigraphical range and show potential to be used, as biostratigraphical indicators through the Eocene sedimentary sequences. The upper part of Gehannam Formation yields vertebrate remains such as the Shark and ray teeth, marine mammal skeletons. Vertebrate remains of great white shark *Macrorhizodus praecursor* (Leriche), *Otodus sokolovi*, *Pristis lathami*, *Pristis aquitinius*, and *Carcharias* sp., in addition to some traces fossils, *Thalassinoides* and *Rhizoliths* are common in Birket Qarun and Qasr El-Sagha Formations. The marked variations in abundance of the recognized vertebrate faunas reflect distinct environmental conditions that control the distribution of many species. The water depth is considered one of the important ecological factors beside the eutrophic state of the water.

The depositional environments in the studied area during Middle-Late Eocene age are interpreted. The abundance of the vertebrate faunas indicates palaeoenvironments varying from open marine shelf with low energy conditions to restricted marine shallow water conditions. However, the abundance of microfauna within calcareous shale and marls of Gehannam Formation indicate an open shallow marine shelf with low energy conditions. The frequent distribution of macrofauna with shallow water shark taxa and intense bioturbation in sandstones of Birket Qarun Formation is

a good indicator of restricted shallow water conditions. Horizontal laminations consisting of alternating shale, siltstone and sandstone with dolostones sheets that show cyclic changes in layer thickness of Temple Member resemble tide-dominated estuaries environment, which cut off by incised valley fill heterolithic sediments of Abu Lifa Member.

References

- Abdallah, A., Abuol Ela, N., Shama, K. and Abdel Aziz, S. (1997). Biostratigraphy of the Temple Member (Qasr El-Sagha Formation, Late Eocene) north El Fayoum depression: Egypt. *J. Geol.* 41(1): 541-578.
- Abu El Ghar, M. S. (2012). Sequence stratigraphy and cyclicity in the Middle Eocene of the Fayoum ranges, Western Desert, Egypt: Implications for regional sea level changes. *Marine and Petroleum Geology*, 29 (1): 276-292.
- Adnet, S. (2006). Nouvelles faunes de sélaciens (Elasmobranchii, Neoselachii) de l'Éocène des Landes (Sud-Ouest, France). Implication dans les connaissances des communautés d'eaux profondes. *Palaeo Ichthyologica*, 10: 1-128
- Adnet, S., Cappetta, H. and Nakaya, K. (2006). Dentition of elasmobranch shark *Mioscyllium* (Squaliformes) with comments on the fossil record of lanternsharks. *Cybium*, 30 (4): 305-312.
- Adnet, S., Antoine, P.O., Hassan Baqri, S.R., Crochet J.-Y., Marivaux L., JL. Welcomme J.-L., and Métais, G. (2007). New tropical carcharhinids (chondrichthyes, Carcharhiniformes) from the late Eocene-early Oligocene of Balochistan, Pakistan: Paleoenvironmental and paleogeographic implications *Journal of Asian Earth Sciences* 30: 303-323.
- Adnet, S., Cappetta, H. and Tabuce, R. (2010). A middle-late Eocene vertebrate fauna (marine fish and mammals) from southwestern Morocco; preliminary report: age and palaeobiogeographical implications. *Geological Magazine*, 147: 860-870.
- Adnet, S., Cappetta, H., Elnahas, S. and Strougo, A. (2011). A new Priabonian Chondrichthyans assemblage from the Western desert, Egypt: Correlation with the Fayoum Oasis. *Journal of African Earth Sciences*, 61: 27-37.
- Allam, A., Shama, K. and Zalat, A. (1991). Biostratigraphy of the Middle Eocene succession at Mishgigah, Wadi El Rayan, Libian Desert, Egypt. *Journal of the African Earth Sciences*; 12(3): 449-9.
- Andreev, P. and Motchurova, N. (2010). Fossil Elasmobranchii and Actinopteryg II. *Bulletin of the Natural History Museum*, 3: 115-129

- Antar, M.S.M. (2011). Paleo-environments of the exposed Eocene Sediments between Wadi El-Hitan and east Siwa in the Egyptian Western Desert based on their faunal content especially the vertebrates. Ph.D Thesis, Zagazig University, Faculty of Science, Geology Department, 257 pp.
- Antunes, M.T.(1978). Faunes ichthyologiques du Neogene superieur d'Angola, leur age, remarques sur le Pliocene marin en Afrique australe. *Ciencias da Terra (UNL)*, Lisboa, 4: 59-90.
- Antunes, M.T. and Jonet, S. (1970). Requins de l'Helvétien supérieur du Tortonien de Lisbonne. *Revista da Faculdade de Ciências da Universidade de Lisboa* 2, C, 16(1): 119-280.
- Arambourg, C. (1952). Les Vertébrés fossiles des gisements de phosphates (Maroc, Algérie, Tunisie). Service Géologique du Maroc, Notes et Mémoires 92: 1-372.
- Bajpai, S. and Thewissen, J.G.M. (2002). Vertebrate fauna from Panandhro lignite field (Lower Eocene), District Kachchh, western India. *Curr. Sci.*, v. 82 (5): 507-509.
- Beadnell, H.J.L.(1905). The topography and geology of the Fayum Province of Egypt. Survey Department of Egypt, Cairo, 101 pp.
- Bourdon, J. and Chandler, R. (2007). The Castle Hayne Formation. Middle Eocene Sharks and Rays of North Carolina. <http://www.elasmo.com>.
- Bown, T.M. and Kraus M. J.(1988). Geology and paleoenvironment of the Oligocene Jebel Qatrani Formation and adjacent rocks, Fayum Depression, Egypt: U.S. Geological Survey Professional Paper, 1452: 1-60.
- Butzer, K.W.(1976). Early Hydraulic Civilisation in Egypt, a Study in Cultural Ecology. The University of Chicago Press, Chicago.
- Cappetta, H. (1970). Les séliaciens du Miocène de la région de Montpellier. *Palaeovertebrata*, mém, ext., 1- 139.
- Cappetta, H. (1976). Selaciens nouveaux du London Clay del'Essex (Ypresien du Bassin de Londres): *Geobios*, 9 (5): 551-574.
- Cappetta, H. (1987). Chondrichthyes II. Mesozoic and Cenozoic Elasmobranchii, in Schults, H.P., Kuhn, O.(eds.), *Handbook of Paleoichthyology*: v.3B, 193pp., Stuttgart, New York: Gustav Fischer Verlag,
- Cappetta, H. (2004). *Handbook of Paleoichthyology* 3B. Chondrichthyes 2 Mesozoic and Cenozoic Elasmobranchii. 193 pp., ISBN 10: 3899370465 ISBN 13: 9783899370461
- Cappetta, H. (2006). Elasmobranchii Post-Triadici, (Index specierum et generum). In: W. Riegraf (ed.), *Fossilium Catalogus, I: Animalia*, v.142, pp. 1-472. Backhuys Publishers, Leiden.
- Cappetta, H., Pfeil, F. and Schmidt-Kittler, N. (2000). New biostratigraphical data on the marine Upper Cretaceous and Palaeogene of Jordan. *Newsletters on Stratigraphy* 38: 81-95.
- Carlsen, A.W. and Cuny, G. (2014). A study of the sharks and rays from the Lillebælt Clay (Early-Middle Eocene) of Denmark, and their palaeoecology. *Bulletin of the Geological Society of Denmark*, Vol. 62: 39-88.
- CASE, G.R. (1981). Late Eocene selachian from south-central Georgia. *Palaeontographica*, Abteilung A176: 52-79.
- Case, G.R. and Borodin, P.D.(2000). Late Eocene selachians from the Irwinton Sand Member of the Barnwell Formation (Jacksonian), WKA mines, Gordon, Wilkinson County, Georgia. *Münchner Geowissenschaftliche Abhandlungen, (A: Geologie und Paläontologie)* 39: 5-16.
- Case, G.R. and Cappetta, H.(1990). The Eocene selachian fauna from the Fayum depression of Egypt. *Palaeontographica*, Abteilung A 212 (1-6): 1-30.
- Case, G.R. and West, R.M.(1991). Geology and paleontology of the Eocene Drazinda shale member of the Kirthar formation, central Western Pakistan. Part II: late Eocene fishes. *Tertiary Research* 12 (3-4): 105-120.
- Case, G.R., Udovichenko, N., Nessow, L., Averianov, A. and Borodin, P.(1996). A Middle Eocene Selachian Fauna from the White Mountain Formation of the Kizylkum Desert, Uzbekistan. *C.I.S. Palaeontographica Abteilung A* 242: 99-126.
- Casier, E. (1949). Contributions à l'étude des poissons fossiles de la Belgique. VIII. Les Pristidés éocènes. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, 25: 1-52.
- Casier, E.(1971). Sur un matériel ichthyologique des "Midra (and Saila) Shales" du Qatar (Golfe Persique). *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique* 47 (2): 1-9.
- Cicimurri, D.J. (2007). A partial rostrum of the sawfish *Pristis lathami* Galeotti, 1837, from the Eocene of South Carolina. *Journal of Paleontology*, 81: 597-601.
- Cicimurri, D.J. and Knight, J.L.(2009). Late Oligocene sharks and rays from the Chandler Bridge Formation, Dorchester County, South Carolina, USA. *Acta Palaeontologica Polonica* 54 (4): 627-647.

- Cione, A.L. and Reguero, M. (1994). New records of the sharks *Isurus* and *Hexanchus* from the Eocene of Seymour Island, Antarctica. *Proceedings of the Geologists Association* 105: 1-14.
- Compagno, L.J.V., Didier, D.A. and Burgess, G.H. (2005). Classification of Chondrichthyan Fish In: Fowler, S.L., Cavanagh, R.D., Camhi, M., Burgess, G.H., Cailliet, G.M., Fordham, S.V., Simpfendorfer, C.A. and Musick, J.A (eds). *Sharks, Rays and Chimaeras: The Status of the Chondrichthyan Fishes. Status Survey. IUCN/SSC Shark Specialist Group*. 461 pp. IUCN, Gland, Switzerland and Cambridge, UK
- Cook, T.D., Murray, A.M., Simons, El. L., Attia, Y.S. and Chatrath, P.(2010). A Miocene selachian fauna from Moghra, Egypt. *Historical Biology*, 22: 1-3: 78-87.
- Dartevelle, E. and Casier, E. (1942). Les poissons fossiles de l'Angola. *Comunicações dos Serviços Geológicos de Portugal*, 22: 99-109.
- Dartevelle, E. and Casier, E. (1943). Les poissons fossiles du Bas-Congo et des régions voisines. *Annales du Musée du Congo Belge, Sér.A (Minéralogie Géologie, Paléontologie)*, 3: 1-200.
- Dartevelle, E. and Casier, E. (1949). Les poissons fossiles du Bas-Congo et des régions voisines. *Annales du Musée du Congo Belge, Sér.A (Minéralogie Géologie, Paléontologie)*, 2: 205-255.
- Dartevelle, E. and Casier, E. (1959). Les poissons fossiles du Bas-Congo et des régions voisines. *Annales du Musée du Congo Belge, Sér. A (Minéralogie Géologie, Paléontologie)*, 2(3): 257-568.
- Diedrich, C.G. (2012). Eocene (Lutetian) Shark-Rich Coastal Paleoenvironments of the Southern North Sea Basin in Europe: Biodiversity of the Marine Fürstenau Formation Including Early White and Megatooth Sharks. *International Journal of Oceanography* Volume 2012: 1-22. doi:10.1155/2012/565326
- Dolson, J., El Barkooky, A., Wehr, F., Gingerich, P.D., Prochazka, N. and Shann, M.(2002). The Eocene and Oligocene Paleogeography and Paleogeography of Whale Valley and the Fayoum Basins: Implications for Hydrocarbon Exploration in the Nile Delta and Eco-Tourism in the Greater Fayoum Basin. Cairo 2002, AAPG/EPEX/ SEG/EGS/EAGE Field trip-guidebook: 1-79.
- Ghosh, B.K. (1959). Some fossil teeth from the tertiary deposits of Orissa. *Jour. Palaeont.*, 33(4): 675-679.
- Gingerich, P.D.(1992). Marine mammals (Cetacea and Sirenia) from the Eocene of Gebel Mokattam and Fayoum, Egypt: stratigraphy, age and paleoenvironments. *Papers on Paleontology* 30: 1-84.
- Hassan, F.A.(1985). Radiocarbon chronology of neolithic and predynastic sites in upper Egypt and the Delta. *African Archaeological Review* 3: 95-116.
- Hassan, F.A.(1986). Holocene lakes and prehistoric settlements of the Western Faiyum. *Egyptian Journal Archaeological Science* 13: 483-501.
- Hassan, F.A.(1997). The dynamics of a riverine civilization: a geoarchaeological perspective on the Nile valley, Egypt. *World Archaeology* 29 (1):51-74.
- Hendrickx, S. and Vermeersch, P.(2000). Prehistory from the Palaeolithic to the Badarian culture. In: Shaw, I. (Ed.), *The Oxford History of Ancient Egypt*. Oxford University Press, Oxford: 17-43.
- Iskander, F. (1943). Geological Survey of the Gharag el-sultany sheet no.68/54: Standard Oil Company, Egypt s. a., reports, 51: p.1-29.
- Ismail, A.S.A. and Abd El-Azeam, S. (2008). Biostratigraphy and paleoenvironment of the Middle-Upper Eocene Ostracods from west Fayoum province-Bulletin of the Faculty of Science Zagazig University; 30: 261-4.
- Itoigawa, J., Nishimoto, H. and Karasawa, H. (1985). Miocene fossils of the Mizunami group, central Japan. 3. Elasmobranchs. *Monograph of the Mizunami Fossil Museum*, 5: 1-89.
- Kemp, D., Kemp, L. and Ward, D. (1990). An illustrated guide to the British Middle Eocene vertebrates, 59 pp. David Ward, London.
- Kent, B.W. (1994). Fossil Sharks of the Chesapeake Region. Egan Rees and Boyer (Eds.): 146 pp. Maryland.
- Kent, B.W. (1999). Part 3. Rays from the Fisher/Sullivan Site. In "Early Eocene vertebrates and plants from the Fisher/Sullivan Site (Nanjemoy Formation) Stanford County, Virginia. Virginia Division of Mineral Resources, Publication, 152: 39-51.
- Kirk, E.C. and Simons, El.L. (2001). Diets of fossil primates from the Fayoum Depression of Egypt: a quantitative analysis of molar shearing. *Journal of Human Evolution* 40: 203-229
- Kozłowski, J.K.(1983). Qasr el-Sagha 1980: Contributions to the Holocene Geology, the Predynastic Settlements in the Northern Fayoum Desert. *Panstwowe Wydawnictwo Naukowe*, Warsaw.
- Kriwet, J. (2005). Additions to the Eocene Selachian fauna of Antarctica with comments on Antarctic

- selachian diversity. *Journal of Vertebrate Paleontology* 25 (1): 1-7.
- Kumar, K. and Loyal, R.S. (1987). Eocene Ichthyofauna from the Subathu Formation, northwestern Himalaya, India. *Jour. Palaeont. Soc. India*, 32: 60-84.
- Leriche M. (1926). Les poissons Néogènes de la Belgique. *Mémoires de Musée Royal du d'Histoire Naturelle de Belgique* 32:367-472.
- Leriche, M. (1927). Les poissons de la Molasse Suisse. *Mémoires de la Société Paléontologique Suisse*, 46: 1-119.
- Leriche, M. (1957). Les poissons neogenes de la Bretagne, de l'Anjou et de la Touraine. *Mem. Geol. Soc. France, N. Ser.*, 36, Mem. 81: 1-61, Paris.
- Longbottom, A.E. (1979). Miocene shark's teeth from Ecuador. *Bulletin of the British Museum natural History, Geology*, 32 (1): 57-70.
- Lowery, D., Godfrey, S.J. and Eshelman, R.(2011). Integrated Geology, Paleontology, and Archaeology: Native American use of Fossil Shark Teeth in the Chesapeake Bay Region. *Archaeology of Eastern North America* 39:93-108.
- Mehrotra, D.K., Mishra, V.P. and Srivastava, S. (1973). Miocene sharks from India. *Rec. Res. Geol. Hindustan Publishing Corporation, Delhi*: 180-187.
- Mondal, S., Das, S. Mallick, S. and Adhikary, D. (2009). Miocene shark teeth assemblages and ancillary fish taxa from baripada, orissa: taxonomic revision and a global Palaeobiogeographic overview. *Journal of the Palaeontological Society of India Volume*, 54(2): 135-152
- Murray, A.M., Cook, T.D., Attia, Y.S., Chatrath, P. and Simons, E.L.L.(2011). A freshwater ichthyofauna from the Late Eocene Birket Qarun Formation, Fayoum, Egypt. *Journal of Vertebrate Paleontology* 30: 665-680.
- Mustafa, H.A. and Zalmout, I.S.(2002). Elasmobranchs from the late Eocene Wadi Esh- Shallala Formation of Qa' Faydat ad Dahikiya, east Jordan. *Tertiary Research* 21: 77-94.
- Mustafa, H.A., Zalmout, I.S., Smadi, A.A. and Nazzal, J.S. (2005). Review of the middle Eocene (Lutetian) selachian fauna of Jebal eth Thuleithuwat, east Jordan. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 237: 399-422.
- Otero, R.A., Oyarzún, J.L., Soto-Acuña, S., Yury-Yáñez, R.E., Pawellek, T. and Adnet, S., Cappetta, H., Metais, E., Salem, M., Brunet, M. and Jaeger, J.-J. (2012). Discovery of an earliest Pliocene relic tropical fish fauna in a newly detected cliff section (Sabratih Basin, NW Libya). *Neues Jahrbuch für Geologie und Palaontologie, Abhandlungen*, 266: 93-114
- Patnaik, R., Sharma, K.M., Mohan, L., Williams, B.A., Kay, R. and Chatrath, P. (2014). Additional Vertebrate Remains from the early Miocene of Kutch, Gujarat. Special publication of the palaeontological society of India, No. 5: 335-351. ISBN: 978-81-926033-2-2
- Priem, F.(1905). Sur les poissons fossiles de l'Eocène moyen d'Egypte. *Bulletin de la Société géologique de France* 5: 633-641.
- Priem, F.(1908). Sur des vertébrés de l'Eocène d'Egypte. *Bulletin de l'Institut Egyptien* 5 (2): 1-3.
- Rana, R.S., Kumar, K. and Singh, H. (2004). Vertebrate fauna from the subsurface Cambay Shale (Lower Eocene), Vastan Lignite Mine, Gujarat, India. *Current Science*, 87 (12): 1726-1733
- Rana, R.S., Kumar, K., Loyal, R.S., Sahni, A., Rose, K.D., Mussell, J., Singh, H. and Kulshreshtha (2006). Selachians from the Early Eocene Kapurdi Formation (Fuller's Earth), Barmer District, Rajasthan, India. *Jour. Geol. Soc. India*, v. 67: 509-522.
- Sahni, A. and Mehrotra, D.K. (1981). Elasmobranch from the coastal Miocene sediments of peninsular India. *Biological Memoirs*, 5 (2): 83-121.
- Sahni, A. and Mishra, V.P.(1975). Lower Tertiary vertebrates from western India. *Monograph of the Paleontological Society of India, Lucknow*, 3: 1-48.
- Said, R. (1962). *The Geology of Egypt*: 377 pp. Elsevier Publishing Co, Amsterdam, New York.
- Said, R. (1990). Cenozoic: in Said, R. (ed.), *The Geology of Egypt*. pp. 451-486, Balkema, Netherlands.
- Said, R., Wendorf, F., Albritton, G., Schild, R., Kobusiewicz, M.(1972). Remarks on the Holocene geology and archaeology of Northern Faiyum Desert. *Archaeologica Polona* 13: 7-22.
- Shama, K. and Shided, A. (1994). Contributions to the Microfiches and Paleogeological Investigation of the Middle and Upper Eocene Sediments in Fayoum area. *Bull. Fac. of Sci., Assiut Univ. Egypt*, 23 (2-f): 381-418.
- Sharma, K.M. and Patnaik, R. (2013). Additional Fossil Batoids (Skates and Rays) from the Miocene Deposits of Baripada Beds, Mayurbhanj District, Orissa, India. *Earth Science India, Vol. 6 (IV)*: 160 - 184.
- Simons, E.L.L. (2005). Eocene and Oligocene Mammals of the Fayoum, Egypt, First International Conference on the Geology of the Tethys, Cairo University, November: 439-450.

- Simons, E.L., Seiffer, E.R., Ryan, T.M. and Attia, Y. (2007). A remarkable female cranium of the early Oligocene anthropoid *Aegyptopithecus zeuxis* (Catarrhini, Propliopithecidae). *National Academy of Sciences of the USA PNAS*. 104 (21): 8731-8736
- Singh, R. (1985). Contribution to the palaeontology and Biostratigraphy of the Subathu Formation of Simla Hills region. Himachal Pradesh. Unpublished Ph.D. thesis, Panjab University: 1-199.
- Smadi, A.A., Abu Azzam, H., Zalmout, I.S. and Mustafa, H.A. (2003). Middle Eocene selachian fauna from Al-Rijla Al-Bayda, Eastern Desert of Jordan. *Abhath Al-Yamouk Basic Science and Engineering*, 12: 619-631.
- Stromer, E. (1905). Die Fischreste des mittlern und oberen Eocäns von Ägypten. I. Teil: Selachii, B. Squaloidei, und II. Teil: Teleostomi, A. Ganoidei. *Beiträge zur Paläontologie und Geologie Österreich-Ungarns und Orients*, Universitäts-Buchhandler in Wien 18: 163-192.
- Stromer, E. (1905a). Die Fischreste des Mittleren und Oberen Eocäns von Ägypten. I. Teil: Selachii, B. Squaloidei, und II. Teil: Teleostomi, A. Ganoidei. *Beiträge zur Paläontologie und Geologie Österreich-Ungarns*, 18: 37-58.
- Stromer, E. (1905b). Die Fischreste des Mittleren und Oberen Eocäns von Ägypten. I. Teil: Die Selachier, A. Myliobatiden und Pristiden. *Beiträge zur Paläontologie und Geologie Österreich-Ungarns*, 18: 163-185.
- Strougo, A. (2008). The Mokattamian stage: 125 years later. *Middle East Research Center. Earth Science Series*, Ain Shams University, vol. 22: 47-108.
- Strougo, A., Cappetta, H. and Elnahas, S. (2007). A remarkable Eocene ichthyofauna from the Elgedida glauconitic sandstone, Bahariya oasis, Egypt, and its stratigraphic implications. *Earth Science Series*, Middle East Research Center, Ain Shams University, vol. 21: 81-98.
- Tiwari, R.P., Mishra, V.P. and Lyngdoh, B.C. (1998). Lower Miocene fish teeth from Mizoram, India. *Geoscience Journal*, 19: 9-17.
- Underwood, C.J., Ward, D.J., King, C., Antar, S.M., Zalmout, I.S. and Gingerich, P.D. (2011). Shark and ray faunas in the middle and late Eocene of the Fayoum area, Egypt. *Proceedings of the Geologists' Association*, 122: 47-66.
- Vliet, H.J. van and Abu El Khair, G. (2010). A new Eocene marine mammal site in the Qattara depression (Egypt). *Cainozoic Research*, 7: 73-77.
- Wallett, L.A. (2006). Eocene selachian fauna from nearshore marine deposits, Ampazony, northwestern Madagascar. Bsc. Thesis, 59 pp. Department of Biological Sciences, South Hadley.
- Weiler, W. (1929). Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wüsten Ägyptens. V. Tertiäre Wirbeltiere. 3. Die Mittel- und obereocäne Fischfauna Ägyptens mit besonderer Berücksichtigung der Teleostomie. *Abhandlungen der Bayerischen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung, neue Folge*, 1: 1-57.
- Welton, B.J. (1972). Fossil sharks in Oregon. *The Ore Bin*, v. 34: 161-170.
- Wendorf, F. and Schild, R. (1976). *Prehistory of the Nile Valley*. Academic Press, New York: 222-226.
- White, E.I. (1926). Eocene fishes from Nigeria. *Bull. Geol. Surv. Nigeria*, v. 10: 1-87.
- White, E.I. (1956). The Eocene fishes of Alabama. *Bulletins of American Paleontology*, 36: 123-151.
- Zalat, A.A. (2015). Holocene diatom assemblages and their palaeoenvironmental interpretations in Fayoum depression, Western Desert, Egypt. *Quaternary International* 369: 86-98.
- Zalmout, I.S.A., Antar, M.S.M., Abd-El Shafy, E., Metwally, M.H., Hatab, E.E. and Gingerich, P.D. (2012). Priabonian sharks and rays (Late Eocene: Neoselachii) from Minqar Tabaghbagh in the western Qattara depression, Egypt. *Museum of Paleontology, University of Michigan*. 32 (6): 71-90.




Figure 1. Geological map of Fayoum depression showing position of the studied stratigraphic sections: I. Hussein Wally, II. Birket Qarun, III. Qasr El Sagha area and IV. Wadi El-Afreet.




Figure 2. Field photograph of A: Hussein Walley section, B: Birket Qarun section, C: Qasr el Sagha section and D: Wadi El-afreet section.




Figure 3. Middle to Late Eocene studied successions A) Hussein Walley village, B) Birket Qarun, C) Qasr El Sagha, D) Wadi El-afreet.




Plate 1: 1-4. *Otodus sokolovi* (Jaekel) 1. lingual view, 3-4. labial views; 5. *Macrorhizodus praecursor* Leriche, lingual view, 6. Fish-fin spine (labial view); 7-8. *Rhizoprionodon* sp., lingual view; 9. *Carcharodon carcharias* Linnaeus, lingual view; 10. *Negaprion frequens* Dames, lingual view; 11,12. *Cretolamna twiggsensis* Case, 11. lingual view, 12. Labial view; 13. *Eutrichiurides* sp., labial view; 14. *Galeocerdo cuvieri* Péron & Lesueur, lingual view; 15. *Hexanchus agassizi* Cappetta, labial view of lower lateral tooth; 16. *Rhinoptera sherburni* Arambourg; 17. *Leidybatis* sp.; 18. *Pristis lathami* Galeotti. (scale bar= 10 mm).

دراسة تصنيفية وتشكلية لبقايا الفقاريات من القروش والاسماك من تتابع عصر الأيوسين الاوسط والعلوى فى منخفض الفيوم ، مصر
 ا.د/ عبد الفتاح على زلط - ا.د/ حمزة مصطفى خليل - د/ محمد صبحى فتحى - رنا محمد طارق
 قسم الجيولوجيا - كلية العلوم - جامعة طنطا

يتناول هذا البحث دراسته الوضع التصنيفي و الشكل الوصفي لبقايا الفقاريات كالقروش والاسماك التي تم تجميعها من تتابع عصر الايوسين الاوسط والعلوى من منخفض الفيوم بالصحراء الغربية مصر من خلال اربعة قطاعات رسوبية تمت دراستها ووصفها وهم على النحو التالي : قطاع قريه حسين والى ، قطاع بركة قارون ، قطاع قصر الصاغة وقطاع وادى العفريت وبناء على الدراسات الطباقية الصخرية لهذه القطاعات فانها تتكون من ثلاث تكوينات مرتبه من الاقدم الى الاحدث كالاتى : متكون جهنم ومتكون بركة قارون ومتكون قصر الصاغة حيث تم التعرف على ٣ فصائل و٧ رتب و١٢ عائلة و ١٨ جنس و ٢١ نوع من أسماك القرش فى منطقه الدراسه وتم وصف الوضع التصنيفي والشكلي لكل نوع والتعرف على البيئات الترسيبية فى عصر الايوسين الاوسط والعلوى فى منطقه الدراسه من خلال بقايا الفقاريات التي تم تجميعها والتي اوضحت ان البيئه الترسيبية تتنوع بين البيئه البحرية العميقة الى البيئه البحرية الضحلة وبالرغم من ذلك فان وجود تنوع من الاحافير الكبيرة واثارها فى الحجر الرملى لمتكون بركة قارون فهذا دليل جيد على البيئه البحريه الضحلة.